



NUI MAYNOOTH

Ollscoil na hÉireann Má Nuad

**OLLSCOIL NA hÉIREANN, MÁ NUAD**

**THE NATIONAL UNIVERSITY OF IRELAND MAYNOOTH**

**First Computer Science and Software Engineering Examination**

**Year 1**

**SEMESTER 2  
2003-2004**

**SE120  
DISCRETE STRUCTURES 2**

Dr. Andrew Martin, Prof. R. Reilly, Mr. T. Naughton

Time allowed: 2 hours

Answer *three* questions

**All questions** carry equal marks

---

1. Prove that the following Hoare triple is correct. Assume that all variables are integer variables. [25 marks]
- ```

{ $x > 0$ }
 $a := 1;$ 
 $b := -1;$ 
while ( $a \leq x$ ) do
   $b := b + 1;$ 
   $a := a * 2;$ 
od
 $a := a/2;$ 
{ $2^b \leq x \wedge 2^{b+1} > x \wedge a = 2^b$ }

```
2. (a) State one predicate that is strictly stronger and one predicate that is strictly weaker than each of the following predicates. You must identify which of your two answers is the stronger and which is the weaker. Furthermore, each of your eight predicates must be unique (you cannot reuse an answer to one part in another part). [8 marks]
- $(x = 5)$
  - $(x = 5 \wedge y = 3)$
  - $(x > 5 \wedge y = 0)$
  - $(x > y \wedge y > 0)$
- (b) For each of the following sets, prove that it is countable or prove that it is uncountable. Note,  $\mathbb{R}_0^1 = \{x : x \in \mathbb{R}, 0 \leq x \leq 1\}$ . [17 marks]
- $\mathbb{N} \times (\mathbb{N} \cup \mathbb{Z})$
  - $\mathbb{N} \times \mathbb{R}_0^1 \times \mathbb{Z}$
  - $2^{\mathbb{N}} \cup \mathbb{N}$
3. (a) Write out three elements of each of the following relations. [9 marks]
- $ABS = \{(a, b) : a \in \mathbb{Z}, b \in \mathbb{N}, \text{abs}(a) = b\}$
  - $SBA = \{(b, a) : a \in \mathbb{Z}, b \in \mathbb{N}, \text{abs}(a) = b\}$
  - $SWAP = \{((a, b), (c, d)) : a, b, c, d \in \mathbb{N}, a = d, b = c\}$
- (b) For each of the relations in part (a) of this question, state whether it is a function or not. For each relation that is not a function, redefine it without changing its meaning so that it is a function. [6 marks]
- (c) Let  $\sim$  be a relation on  $B = \{0, 1, 2, 5, 7, 11, 12\}$  defined by  $x \sim y$  iff  $\text{mod}(x, 3) = \text{mod}(y, 3)$ . Use this relation to partition  $B$ . [5 marks]
- (d) Given the binary relation  $R = \{(0, 1), (1, 2), (2, 3), (4, 4)\}$  construct each of the following compositions. [5 marks]
- $R^2$
  - $R^3$
  - $R^4$

4. (a) Each of the following Hoare triples claims to correctly perform the swapping process. The first one uses a temporary variable. The second does not. Prove that each triple is correct. Assume that all variables are integer variables. [15 marks]
- $\{x < y\} \text{ temp} := x; x := y; y := \text{temp} \{y < x\}$
  - $\{x < y\} y := y + x; x := y - x; y := y - x \{y < x\}$
- (b) Let the language  $L = \{aa, ab, bbb, aaab, abab, baaa\}$ . Define an equivalence relation of your choice that partitions  $L$  into the following numbers of classes. [10 marks]  
Write out the contents of the classes in each case.
- An equivalence relation that partitions  $L$  into exactly two classes
  - An equivalence relation that partitions  $L$  into exactly four classes

### Rules that can be applied in any question

|                          | $P$ | $Q$ | $P \rightarrow Q$ |
|--------------------------|-----|-----|-------------------|
| Implication truth table: | T   | T   | T                 |
|                          | T   | F   | F                 |
|                          | F   | T   | T                 |
|                          | F   | F   | T                 |

Simplification (Simp):  $\frac{A \wedge B}{A}$

Addition (Add):  $\frac{A}{A \vee B}$

Conjunction (Conj):  $\frac{A, B}{A \wedge B}$

Transitive:  $\frac{a > b \wedge b > c}{a > c}$

Conditional Proof Rule (CP): If there is a proof of  $B$  from the assumption that  $A$  is true (i.e. if  $B$  can be derived from  $A$ ), then  $A \rightarrow B$

Assignment Axiom (AA):  $\{Q(x/t)\} x := t \{Q\}$

Consequence Rule:  $\frac{P \rightarrow R \quad \text{and} \quad \{R\} S \{Q\}}{\{P\} S \{Q\}}$

Composition Rule: 
$$\frac{\{P\} S_1 \{R\} \quad \text{and} \quad \{R\} S_2 \{Q\}}{\{P\} S_1; S_2 \{Q\}}$$

If-Then Rule: 
$$\frac{\{P \wedge C\} S \{Q\} \quad \text{and} \quad P \wedge \neg C \rightarrow Q}{\{P\} \text{ if } C \text{ then } S \{Q\}}$$

If-Then-Else Rule: 
$$\frac{\{P \wedge C\} S_1 \{Q\} \quad \text{and} \quad \{P \wedge \neg C\} S_2 \{Q\}}{\{P\} \text{ if } C \text{ then } S_1 \text{ else } S_2 \{Q\}}$$

While Rule: 
$$\frac{\{P \wedge C\} S \{P\}}{\{P\} \text{ while } C \text{ do } S \{P \wedge \neg C\}}$$

Statements that can be quoted without proof:

1.  $\mathbb{N}$  is countable
2. Any set that has a bijection with a subset of  $\mathbb{N}$  is countable
3. Let  $B = A_1 \cup A_2 \cup \dots \cup A_n$ . If each  $A_i$  is countable then  $B$  is countable. If at least one  $A_i$  is uncountable then  $B$  is uncountable.
4. Let  $B = A_1 \times A_2 \times \dots \times A_n$ . If each  $A_i$  is countable then  $B$  is countable. If at least one  $A_i$  is uncountable then  $B$  is uncountable.